



COMPREHENSIVE PAIN MANAGEMENT SPECIALISTS

P.O. Box 501724, San Diego, CA 92150
Phone 858.453.7700 Fax 858.798.1225

Constipation Avoidance

Constipation is defined as infrequent bowel movements of the passage of hard stools, causing discomfort.

Steps to prevent constipation: Start at number 1 and move down the numerical steps until bowel movements are soft and regular.

1. Drink 8-10 glasses of fluid a day, including fruit juices. Eat a high fiber diet including raw fruits and vegetables, dried fruits, and whole grains. In addition, try the juice recipe below.

Juice recipe:

- 1 oz prune juice
- 1 oz apricot nectar
- 1 oz water
- 1 oz Karo syrup

Mix in a pitcher and drink 4 oz. every day. Adjust to softness and frequency of stool.

Cereals such as Quaker Oatmeal Squares are high in fiber.

2. Use a STOOL SOFTENER: (Dioctyl sodium sulfosuccinate, DSS), DUCOSATE sodium 250 mg or 1 capsule orally; or Colace 100 mg by mouth 2 times per day. May be purchased without a prescription. Remember to keep up your fluids with juices and water as in Step 1.
3. Take a LAXATIVE/STIMULANT if you have no bowel movement for 3 days. Laxatives increase bowel activity so that stool moves down and out. Take Senokot 2 tablets at night. This may be increased to 2 tablets three times per day as needed (6 tablets per day max).
4. If Senokot 2-6 tabs per day did not work, ass Dulcolax Suppository in the rectum the next morning (Caution: These stronger laxatives can cause cramping).
5. If Dulcolax doesn't result in a bowel movement, take Milk of Magnesia (MOM) 15-30 mls or Lactulose 30 mls (2 tsp). If you have no bowel movement within three hours, take another Dulcolax suppository. If no results by evening take a fleet enema.
6. If this plan does not work or you are having increasing pain or cramping in your abdomen, call your physician.